

Received Date : 25-May-2015

Revised Date : 04-Sep-2015

Accepted Date : 19-Sep-2015

Article type : Perspective

Harnessing natural selection to tackle the problem of prey naïveté

Katherine E. Moseby^{1,2}, Daniel T. Blumstein³, Mike Letnic⁴

¹School of Earth and Environmental Sciences, The University of Adelaide, Adelaide, SA 5005, Australia

²Arid Recovery Ltd., P.O. Box 147, Roxby Downs 5725, Australia

³Department of Ecology and Evolutionary Biology, University of California, 621 Young Drive South, Los Angeles, CA 90095-1606, USA

⁴Centre for Ecosystem Science, School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, NSW 2052, Australia

Katherine.moseby@adelaide.edu.au

M.letnic@unsw.edu.au

marmots@ucla.edu

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/eva.12332

This article is protected by copyright. All rights reserved.

Running head: natural selection and prey naïveté

Abstract

Many populations are threatened or endangered because of excessive predation resulting from individuals' inability to recognize, avoid, or escape alien predators. Such prey naïveté is often attributed to the absence of prior experience and co-evolution between native prey and introduced predators. Many reintroduction programs focus on reducing predation rate by excluding introduced predators, a focus which ignores, and indeed exacerbates, the problem of prey naïveté. We argue for a new paradigm in reintroduction biology that expands the focus from predator control to kick-starting learning and evolutionary processes between alien predators and reintroduced prey. By exposing reintroduced prey to carefully controlled levels of alien predators, *in-situ* predation could enhance reintroduction success by facilitating acquisition of learned anti-predator responses and through natural selection for appropriate anti-predator traits. This *in situ* predator exposure should be viewed as a long term process but is likely to be the most efficient and expedient way to improve prey responses and assist in broadscale recovery of threatened species.

Keywords: exotic predators, prey, naïveté, natural selection, predator learning, reintroduction

The problem of prey naïveté

Predation by introduced predators, especially mammalian predators, is a major factor responsible for the extinction of wild vertebrate populations and the failure to successfully reintroduce endangered vertebrates in many parts of the world (King 1984; Savidge 1987; Biggins et al. 1999; Johnson 2006; Moseby et al. 2011). The primary reasons for significant population declines of many native prey species and failure of subsequent reintroduction programs are thought to be 1) the inability of prey individuals to avoid and or mount effective anti-predator responses when they encounter introduced predators (Griffin, Blumstein, and Evans 2000; Short, Kinnear, and Robley 2002; Blumstein 2006; Moseby et al. 2011), and 2) the high densities and therefore high rates of encounters between introduced predators and native prey.

In many situations introduced predators thrive in their new environments and may occur at very high population densities. There are several potential drivers of high population densities of introduced predators. These include release from constraints on population growth posed by larger predators, competitors and parasites as well as facilitation that can occur when populations of introduced predators benefit from the presence of high densities of introduced or native prey (MacDonald and Harrington 2003; Saunders, Gentle, and Dickman 2010; Sih et al. 2010; Letnic, Ritchie, and Dickman 2012). High rates of encounters between prey and over-abundant predators can have catastrophic effects on prey populations. These population-level effects of predators on prey are likely to be exacerbated and result in “hyper-predation” on rare prey species if alternate food sources (such as over-abundant prey) are available for predators (Sinclair et al. 1998).

While there is evidence that high densities of introduced predators can precipitate catastrophic declines in prey populations and thwart reintroduction attempts, in many instances just one or relatively few individuals of an introduced predator species have had catastrophic impacts on populations of threatened prey (Christensen and Burrows 1994; Gibson et al. 1994; Moseby et al. 2011). These observations suggest that the susceptibility of some native prey to introduced predators is not just a function of the rate of encounters between predators and prey but also due to an asymmetry in the outcome of predator-prey encounters. For example, even when intensive feral animal control was successfully implemented at a site in inland Australia, individual exotic feral cats (*Felis catus*) were still able to cause the failure of a large-scale mammal reintroduction program (Christensen and Burrows 1994). Similarly, Roy et al. (2002) reported that population control of an introduced mongoose (*Herpestes javanicus*) failed to prevent uncommon but significant predation events on the rare pink pigeon (*Colombus mayeri*) in Mauritius.

The high susceptibility to introduced predators displayed by many prey species is at least partly due to their naïveté (Sih et al. 2010; Carthey and Banks 2014). Naïveté can result from isolation of individuals from predators during their lifetime (ontogenetic naïveté, common in captive-bred animals) or, through an absence of co-evolution with recently introduced predators (evolutionary naïveté, common in situations where exotic predators are present) (Griffin, Blumstein, and Evans 2000). Banks and Dickman (2007) suggested that there are three levels of naïveté to alien predators: level 1 is a failure of prey to recognize a species as a predator; level 2 is recognition of the predator but adoption of inappropriate anti-predator behavior; and level 3 is where prey recognize the predator, have an appropriate response, but the predator exhibits

superior hunting skills. In addition to inappropriate anti-predator behavior, prey species that lack evolutionary exposure to predators may also possess other traits that make them susceptible to novel predators such as flightlessness, strong scent, noisy or conspicuous young, inadequate camouflage and lack of nest or brood guarding behaviors.

Prey naïveté has been a particularly problematic issue for reintroduction programs in Australasia. Here, vulnerability to predation by introduced mammalian predators has caused widespread declines and extinction of more than 20 species of mammals weighing less than 5 kg in Australia (Johnson 2006) and numerous species of birds and lizards in New Zealand (King 1984; Towns, Daugherty, and Cree 2001). In some cases, populations of endangered native species remain in areas where introduced predator populations are low or absent (Johnson 2006; Innes et al. 2010). These remnant populations have been used as sources for reintroduction programs into areas of their former ranges where introduced predators exist, but in nearly all cases reintroduction programs have failed to establish self-sustaining wild populations (Christensen and Burrows 1994; Short and Turner 2000; Moseby et al. 2011; Hayward et al. 2012).

Predation from introduced predators has in most cases been cited as the primary cause of reintroduction failure in Australasia (Fischer and Lindenmayer 2000; Short and Turner 2000; Short 2009; Sherley et al. 2010; Moseby et al. 2011). In Australia, predation by introduced predators, the red fox (*Vulpes vulpes*) and feral cat was the reason for failure of approximately 80% of unsuccessful mammal introductions (Short 2009). In taxa such as macropodids (kangaroos, wallabies and rat kangaroos) their susceptibility to introduced predators is so great

that no safe density of introduced predators is thought to exist (Clayton et al. 2014). However, native predators can also cause reintroduction failure, particularly in captive-bred animals, with predation and prey naivety cited as primary reasons for the failure of bird reintroductions in the Caribbean (White Jr, Collazo, and Vilella 2005) and Saudi Arabia (van Heezik, Seddon, and Maloney 1999).

In this perspective, we propose a novel approach to overcome the problem of prey naïveté that aims to kick-start co-evolution between introduced predators and native prey. While much of our discussion is focused on improving the success of endangered species reintroduction programs, the concepts and approaches that we outline have broader relevance and application to ecosystems where native prey interact with introduced predators.

Expanding the focus of reintroduction programs from predators to prey

To date, strategies to address high predation rates in reintroduction programs have largely concentrated on controlling or eradicating predators rather than improving the anti-predator responses of prey (Armstrong et al. 2002; Scofield, Cullen, and Wang 2011). However, with the exception of some island or fenced systems where a well-defined area and the absence of immigration can facilitate predator eradication (Nogales et al. 2004), the likelihood of eradicating introduced predators is low. Consequently, many reintroduction programs focus on improving the survival of reintroduced species by reducing predator population sizes and thereby reducing the frequency of encounters between threatened prey and predators. For example, in Australia and New Zealand, one widely employed strategy to control introduced predators (primarily red foxes in Australia, and stoats (*Mustelus ermina*) and rats (*Rattus rattus*) in New

Zealand) is the broad scale distribution of poisoned baits, often from aircraft (Burrows et al. 2003; Moseby and Hill 2011; Ruscoe et al. 2011). Another strategy that has been advocated to alleviate introduced predators impacts on prey, but remains untested in reintroduction programs, is to harness the suppressive effects that native apex predators have on populations of introduced predators (most introduced predators are mesopredators) (Crooks and Soulé 1999; Letnic, Ritchie, and Dickman 2012; Hunter et al. 2015).

In larger areas, strategies for endangered species persistence that rely on suppressing or excluding populations of introduced predators may only be effective in the short-term (Wayne et al. 2013). There are at least two reasons why lasting predator control is difficult to achieve. First, control techniques such as poisoning, biological control, and shooting/trapping may impose selection for predators that are less susceptible to the control technique and thus become less effective over time (Warburton and Drew 1994; Allen, Engeman, and Krupa 1996; Kohn, Pelz, and Wayne 2000). Second, programs that require perennial support from funding sources to implement predator control are vulnerable to factors beyond the control of land-managers, such as variation in financial markets and the whims of funding agencies and philanthropists. Any of these reasons could lead to the failure of a reintroduction program if introduced predator populations cannot be suppressed to sufficiently low levels to allow the persistence of reintroduced species. Thus, while control of introduced predators can assist in the protection of threatened species populations (see Marlow et al. 2015), results can be short-lived (Cote and Sutherland 1997) and there is a need to explore other alternatives for long term co-existence.

In addition to evolutionary naivety, another problem facing threatened species reintroduction programs is ontogenetic naivety. Necessity and availability dictates that threatened species reintroduction programs often use captive-bred stock or animals sourced from introduced predator-free islands or fenced enclosures (van Heezik, Seddon, and Maloney 1999; Griffin, Blumstein, and Evans 2000; Short and Turner 2000). Indeed, in Australasia, the majority of threatened species reintroductions now involve the transfer of individuals between predator-free refuges (Towns and Ferreira 2001; Short 2009; Scofield, Cullen, and Wang 2011). Although threatened prey-species are kept safe from predators and reintroduction success is inevitably improved, the problem of prey naïveté is exacerbated by predator exclusion because the predator avoidance strategies of individuals from such predator-free areas are often severely compromised (Biggins et al. 1999; McLean et al. 2000).

Complete removal of predation pressure can lead to significant and rapid loss of prey anti-predator behavior through relaxed selection (Blumstein 2006; Lahti et al. 2009). This is because anti-predator behaviors often involve animals increasing their anti-predator vigilance at the expense of engaging in other activities, or animals avoiding certain areas because there may be an increased risk of predation. Thus, with no predators, there are no benefits from engaging in anti-predator behavior and it may be rapidly lost (Blumstein, Daniel, and Springett 2004). The often poor anti-predator response of prey from predator-free sanctuaries can hamper efforts to reintroduce individuals from these small, isolated systems into larger, natural areas in their former ranges where predators now exist.

The multi-predator hypothesis (Blumstein 2006) states that prey should be retain the ability to respond to predators, even extinct ones, as long as they are exposed to some predators. This is because whenever prey have more than a single predator, we expect that anti-predator behaviors will not assort independently and we expect the evolution of anti-predator syndromes. Thus, populations exposed to some predators, even those from different archetypes (e.g., avian but not mammalian), may retain predator discrimination abilities for all archetypes (avian and mammalian) and effective anti-predator behavior even for the missing predator(s). Support for the multi-predator hypothesis comes from studies that document long-term persistence of anti-predator behavior despite the loss of key predators (Byers 1997; Blumstein, Daniel, and Springett 2004). Thus, some predation may be essential to retain the ability to respond to predators. True isolation in completely predator-free enclosures may, however, lead to a rapid loss of anti-predator abilities.

The predator archetype hypothesis predicts that if two predators are similar in some key way, species will respond to them (Cox and Lima 2006). For instance, there is remarkable morphological convergence among mammals that have similar diets and hunting styles (Wroe and Milne 2007). Additionally, related species may share similar olfactory chemicals. Support for the olfactory archetype hypothesis comes from studies of fishes trained to recognize one predator are more likely to generalize towards more closely-related predators and less likely to generalize towards more distantly related predators (Ferrari et al. 2007). Therefore, by addressing the problem of prey naïveté rather than trying to eliminate all predators, the risks to prey from future encounters with predators may be reduced.

Another problem associated with complete removal of predators from ecosystems can arise if populations of reintroduced species increase to a point where their consumption of resources has adverse effects on the ecosystems to which they have been introduced (Hayward and Kerley 2009). Overgrazing by herbivore populations that are unchecked by natural predators has been linked to environmental degradation and fluctuations in herbivore populations in unfenced ecosystems around the world (Côté et al. 2004; Letnic, Ritchie, and Dickman 2012).

Overpopulation is an important consideration in predator-free sanctuaries, because in addition to the environmental damage that can occur, there is a serious risk of catastrophic population declines of reintroduced species and adverse impacts on other species within the sanctuaries if the food resource base is exhausted (Wiseman, Page, and O'Connor 2004; Slotow et al. 2005; Crisp and Moseby 2010; Islam, Ismail, and Boug 2010). In some cases supplementary feeding and or culling is required to manage populations reintroduced into predator-free sanctuaries after animals have over-eaten their resource base (Hayward et al. 2007; Crisp and Moseby 2010).

These situations are clearly not natural and result in the creation of sanctuaries that are more like zoos than ecosystems (Scofield, Cullen, and Wang 2011). While introductions to islands or fenced areas may preserve endangered species, we suggest that if ecological restoration and the establishment of self-sustaining populations of reintroduced species is one of the aims of reintroduction programs then we must move beyond predator-free sanctuaries and create sustainable ecosystems. To accomplish this we must address the inability of prey-species and introduced predators to co-exist.

Given the potential problems associated with achieving sustained control or eradication of introduced predators we argue that there is an urgent need to expand the focus of predator management in reintroduction programs to not just focus on predator removal (which reduces the frequency of encounters between predators and prey), but to also improve the ability of prey species to avoid fatal encounters with introduced predators. While we recognize that predator detection is but one of several stages of the predation process, we suggest that if the anti-predator responses of endangered prey populations could be improved, it might be more likely that introduced predators and endangered prey will co-exist in the wild. An additional benefit of improved anti-predator responses is that some tend to be broad ranging and not necessarily species-specific (Blumstein 2006; Cox and Lima 2006).

Why predator avoidance training involving simulated encounters with predators in captivity is likely not the answer

Lack of predator recognition either through evolutionary or ontogenetic isolation is thought to be the most damaging form of prey naïveté (Cox and Lima 2006). Some reintroduction programs have attempted to improve predator recognition prior to re-introductions by simulating encounters with predators, whereby predator cues are paired with an unpleasant experience (Griffin, Blumstein, and Evans 2000). Such pre-release predator training can modify prey behavior (Miller et al. 1990; Maloney and McLean 1995; Hölzer, Bergmann, and McLean 1995; McLean, Lundie-Jenkins, and Jarman 1996) and empirical evidence shows that fish (Brown and Laland 2003), birds (McLean, Hölzer, and Studholme 1999), and mammals (McLean et al. 2000) can be trained to improve their anti-predator skills. In some cases, trained individuals may serve as demonstrators from which other animals learn to improve their anti-predator skills (Griffin

and Evans 2003; Griffin 2004). If widespread, social transmission of anti-predator behavior could be an effective mechanism by which animals can learn from others' experiences without direct human interventions.

Despite the promise of pre-release training coupled with natural social transmission, only a few studies have empirically tested and demonstrated that pre-release predator training can improve post-release survival of reintroduced species (White Jr, Collazo, and Vilella 2005; van Heezik, Seddon, and Maloney 1999). Indeed, most practitioners investigating the utility of pre-release predator training have used evidence of a pre-release change in behavior of trained individuals rather than a difference in post-release survival of trained and untrained individuals as a measure of success (Miller et al. 1990; Holzer et al. 1996; McLean, Lundie-Jenkins, and Jarman 1996; Moseby, Cameron, and Crisp 2012). Of particular note is that studies reporting improved survival of trained captive-bred animals after release invariably involve a response to native predators rather than to exotic species (White Jr, Collazo, and Vilella 2005; Gaudioso et al 2011; Carthey and Banks 2014). To our knowledge, pre-release training has not been shown to reduce post-release survival in prey species exposed to exotic predators.

In many cases, lab-based predator avoidance training has been unsuccessful because captive situations do not provide the conditioning necessary for survival in the wild. Pre-release predator training often focuses on captive-bred animals that are trained in captivity (Beck et al. 1994). Captive-bred animals tend to have a much lower survival rate than wild animals upon release for a variety of reasons including that individuals are unfamiliar with the release site, can travel large distances, and often exhibit abnormal behavior post-release (Snyder et al. 1996).

Additionally, predator avoidance trials in captivity rarely use real predators but instead have primarily used harassment with stimuli such as with rubber bands, water pistols, stuffed animals mounted on wheels, and loud noises (McLean et al. 2000). Such unrealistic stimuli are unlikely to stimulate the fear conditioning that likely occurs naturally when an animal survives a real predatory encounter (Schakner and Blumstein 2016). Notably, one successful pre-release predator training program which did result in improved post-release survival involved the exposure of houbara bustards (*Chlamydotis undulata*) to live predators prior to their release (van Heezik, Seddon, and Maloney 1999).

Kick-starting learning and natural selection

We suggest that a realistic and potentially useful form of predator avoidance training involves *in situ* predator exposure using real encounters between wild prey populations and predators. There are at least three advantages of exposing prey species to real predators. First, there may be an increased capacity for learning and reinforcement. Second, there is the opportunity to select for individuals with improved anti-predator behaviors through natural selection. Third, predators not humans will do the selecting and thus avoid biases towards particular traits that can be readily observed or quantified by humans.

Learning is improved because the stimuli are real, the exposed population is wild, and cultural transmission can occur during all life stages. Several studies have reported improved learning of anti-predator behavior in captive situations when actual live predators are used rather than predator models or scent (White Jr, Collazo, and Vilella 2005; Carthey and Banks 2014). Additionally, appropriate predator avoidance behavior is likely to be strongly reinforced during

in situ predator exposure due to prolonged exposure to the predator and the opportunities for filial and cultural transfer (Griffin et al. 2000).

Perhaps more importantly, *in situ* predator exposure allows us to harness the effects of natural selection to select for appropriate traits. We envisage, that selection by predators is unlikely to operate on a single trait, but on a suite of characters that could conceivably include behavioral, physiological and physical traits. Thus, we propose that strong selection pressure imposed by introduced predators on some naïve prey species should improve their capacity to avoid fatal encounters with predators. However, it is important to note that the traits linked to improved survival which predators select for may not be readily observable or quantifiable by humans except as improved survival. Thus, an advantage of using *in situ* predator exposure may be that it could reduce biases towards selection for readily measurable traits imposed by humans and instead select for the traits that demonstrably confer greater longevity and reproductive success.

In the case of behavioral responses to predators, the relative significance of learning vs. natural selection in improving anti-predator behavior is likely to be influenced by the sociality of the prey species. Solitary species may rely more on natural selection and filial transfer, while social species may have improved opportunities for learning from conspecifics (Griffin et al. 2000). We caution, however, that learning or enhancing innate anti-predator responses to native predators lost through ontogenetic isolation is likely to be substantially easier than developing new, effective anti-predator responses against introduced predators due to evolutionary naïveté.

As evidenced by the extinction of predator-exposed populations of naïve prey, a major obstacle to overcome when using *in situ* predator exposure would be the magnitude of the selection pressure that predators can impose on naïve prey. A solution may be to expose prey to predators under tightly controlled conditions, where the rate of prey mortality can be closely monitored, and the predators removed should there be a risk that the prey population could become extinct..

From a viewpoint focused on maintaining genetic diversity, it may appear risky to select threatened species for a suite of traits associated with anti-predator responses, because such selection could reduce the population's genetic diversity. However, this initial loss of genetic diversity is likely to be offset if improved learning and the long term co-existence of prey species with exotic predators eventually leads to larger populations of threatened species and, ultimately, greater genetic variation. We suggest that if prey populations are to survive with exotic predators in the long term, then the genetic bottlenecking that inevitably results from a bout of natural selection will be desirable if it improves threatened species' capacity to co-exist with introduced predators. Preserving genetic diversity in captive breeding programs requires significant time and effort. Most diversity is lost when animals are released and high mortality from predation occurs. In the case where prey naïveté and predation from exotic predators is the most important factor causing population extinction, we consider selecting for traits associated with enhanced survival when confronted with introduced predators to be both justifiable and essential: some reduction in genetic variation is better than the complete failure to recover a population in the wild.

Accepted Article

Are there populations of threatened prey species large enough to conduct *in situ* predator exposure? Predator-free islands and fenced sanctuaries offer the best opportunities to manage this *in situ* process. In some cases these areas experience an oversupply of threatened prey species due, ironically, to an absence of natural predators. Sterilized, and radio-collared single predators could be added to islands and fenced enclosures where healthy populations of threatened prey are present whilst leaving adjacent areas untouched as insurance populations. We assume that predators will explore this environment and leave signs of their presence (e.g., scent marks, vocalizations) that prey can learn to associate and potentially avoid. We assume that some prey will be attacked, and some may be killed. Importantly, we assume that some prey will directly survive encounters with predators while others might have the opportunity to learn vicariously through others' experiences. These are realistic assumptions. Many prey species identify and respond to the scents and sounds of their predators; even novel predators that resemble natural predators (Kohn, Pelz, and Wayne 2000; Blumstein et al. 2002; Blumstein 2006; Anson and Dickman 2013; Gérard et al. 2014). Previously naïve survivors have been shown to modify their behavior after direct or vicarious experiences with predators (Berger, Swenson, and Persson 2001). Additionally, previous single predator incursions have been recorded in fenced reserves and have not resulted in mass predation events, suggesting that many prey species can tolerate the low densities of predators required for these trials (see examples in Moseby et al. 2015).

Given these assumptions, prey that survive will either have learned to avoid predation, or undergone a bout of selection through which survivors passed on whatever heritable traits that facilitated survival to the next generation. There are many examples of environmental change

associated with human activities and biological invasions imposing strong selection pressures on species and in doing so driving rapid phenotypic change (Phillips and Shine 2004; Darimont et al. 2009). Such changes can be both evolutionary and/or represent phenotypic plasticity (Kohn, Pelz, and Wayne 2000; Hendry, Farrugia, and Kinnison 2008). The capacity of native species to respond to *in situ* training is likely to vary considerably depending on vulnerability of each life stage to predation and the opportunities that exist for filial and cultural transfer. We suggest that it is essential to understand the life history strategies a species uses as well as documenting baseline naiveté to identify potential responsive populations and species for such *in situ* training.

How do we predict if and when natural selection by predators will sufficiently reduce prey vulnerability in existing predator environments? This can be tested through both intergenerational comparisons and experimental manipulations. The changes in intergenerational prey responses can be measured by quantifying how prey respond to predator cues using scents, sounds and models as well as how their vigilance and escape behavior (e.g., flight initiation distance) varies (Cooper, Diogo, and Blumstein 2015). Additionally, the survival rates of successive generations can be tested during *in situ* predation experiments: increased survival demonstrates improved anti-predator responses.

When differences are detected, manipulations can be conducted to determine when prey are “ready” to face environments that support predator populations. These can be conducted by gradually adding more *in situ* predators up to the densities recorded in existing predator environments and measuring the responses of prey. Ultimately, successive generations of *in situ* trained prey would be released into existing predator areas with varying predator densities and

their survival compared with untrained individuals. Understanding the predation thresholds that can be tolerated by untrained and progressively trained prey will also assist with understanding the limitations of *in situ* predator training and the importance of simultaneous predator control.

An important consideration for programs that use *in situ* predation to improve reintroduced species anti-predator responses will be the threshold level of population reduction at which the experiment is ceased. Theory predicts that the stronger the selection (i.e., the more animals killed), the more likely the resulting population will be different from the original population in its anti-predator behavior. Massive reductions of a threatened or endangered species, however, may not be practical or desirable. In practice, the threshold level of population reduction using *in-situ* predation will be determined by trade-offs between factors such as the number of individuals of the reintroduced species population, the expected rates of intrinsic growth and natural mortality, and the capacity to remove the predator. Such demographic factors could be modelled prior to initiating an experiment.

Ethical considerations

Is this *in situ* predator exposure ethically defensible? Will it result in excessive mortality of endangered species? What about the welfare of individual prey species and predators?

We adopt here an explicitly ecologicistic perspective (e.g., Kellert 1997; Simaika and Samways 2010) because our primary goal as conservation biologists and ecologists is to help recover populations and restore ecosystems. By contrast, Vucetich and Nelson (2007) emphasize the importance of thinking about the welfare of individuals that may suffer in a conservation

intervention. While our focus on population recovery does not mean that we should not be concerned with the welfare of individuals, it does focus our goals on the ultimate goal of recovering populations of threatened or endangered species. We believe that wildlife managers need more effective tools and approaches to recover populations; particularly those where failure is in response to predation on recently released animals.

We believe that the lessons from reintroduction biology have shown us that captive bred animals have a particularly high mortality rate upon release into the wild and that this is often driven by predation. In cases where naïve prey encounter introduced predators, predation may be absolute. Thus, some have argued that reintroductions, given their low success rate, are ethically questionable (Bekoff 2002). However, increasing populations of endangered species and creating sustainable wild populations in some jurisdictions are mandated by laws such as the Endangered Species Act of the USA. If we aim to restore ecosystems where introduced predators interact with naïve prey, something bold must be done. Viewed this way, there is a moral imperative to explore novel management strategies.

Does this need to develop more effective introduction strategies outweigh the costs to individual prey and predators that might be involved in such an intervention? We believe it could and therefore it is certainly worth exploring the idea with a set of well-designed experiments. For our proposed interventions to work, some individual prey will likely die. While we understand that not all will agree with our viewpoint, we believe that if *in situ* predation improves the anti-predator responses of endangered prey, and populations are ultimately recovered, the individual suffering imposed on individual prey who might be killed while in an enclosure with a captive

predator is outweighed by the benefits of recovering the population by programs which aim to reduce the rate of encounters between predators and prey by killing predators.

Issues relevant to the welfare of the predator are worthy of discussion and debate. We envision two types of predators that would be used in this sort of experiment: evolutionarily novel ones and native predators. Neither directly benefits from *in situ* predator exposure; indeed, they are used as tools for its implementation.

Consider introduced cats and foxes in Australia. Introduced by Europeans, these species have had a disproportionately negative impact on small and mid-sized Australian mammals (Johnson 2006). Thus, in many places these introduced predators are primarily poisoned and to a lesser degree trapped and shot to reduce population density or eliminate them in fenced reserves (Moseby et al. 2011). Nonetheless, native Australian mammals must interact with them if they are to survive outside a fenced reserve. Thus, it may be defensible to use these abundant predators, who otherwise would be killed, in experiments to determine the efficacy of *in situ* predator exposure if their welfare is taken into account. Furthermore, if *in situ* predation is successful at improving the anti-responses of endangered mammals it may negate the need to kill introduced predators in the future and so reduce the suffering experienced by predators. By contrast, consider a native predator and a native prey. Depending on the situation, it might warrant further discussion and debate about whether to use a native canid, dingoes in the Australian example, as a tool to prepare a small mammal for release. We think that these are questions that must be discussed and debated on a case-by-case basis (e.g., Vucetich and Nelson 2007).

The ultimate goal of *in situ* predator exposure is to reduce mortality rates so that the rate of predator-driven mortality does not exceed the rate of population growth. Recent studies have called for consideration of both the ecological and evolutionary cost of resource management decisions (Ashley et al. 2003), emphasizing that evolutionary changes can occur over relatively short time frames (Thompson 1998; Hendry, Farrugia, and Kinnison 2008). For instance, managers already recognize that genetic changes can occur over short time frames in animals subjected to captive housing as they adapt to captive conditions (Williams and Hoffman 2009). Recommendations for improving reintroduction success using captive-bred animals include minimizing the number of generations in captivity (Williams and Hoffman 2009). Closer integration of evolutionary biology with reintroduction biology is likely to improve environmental outcomes through what Carroll et al. (2014 pg. 1) term "manipulating the relationships between the traits of organisms and the patterns of selection imposed by their environments". Our suggestion of using *in situ* predation to force rapid behavioral and potentially genetic change is one example of how these manipulations and integrations could be used to improve reintroduction success. We do not envision this technique being possible to adopt in all cases where we have threatened populations of captive animals, but in those cases where it is possible, we think that it may ultimately be a useful tool that might even result in a reduced need to kill extant predators.

A paradigm shift

To date there has been little focus on the role of prey naïveté in the decline of threatened wildlife species despite introduced predators being a prominent cause of species declines. Introductions of the red fox to California and subsequent fauna declines (Lewis, Sallee, and Golightly Jr 1999)

show that the impacts of introduced predators are not confined to island ecosystems or specific countries. The global risk of future predator incursions is high, fuelled by the deliberate and inadvertent movements of species by humans. Reviews have highlighted the importance of advancing our understanding of prey naïveté (Carthey and Banks 2014) particularly in relation to introduced predators. We call for novel strategies to help wildlife biologists protect and re-establish populations of endangered wildlife and strongly encourage a shift in focus from exclusion of introduced predators to improving prey responses in order to facilitate future resilience and possible co-existence. This paradigm shift from predator exclusion to co-existence would not diminish the importance of predator control, but rather assist in facilitating an integrated, multi-level approach to threatened species management.

The effectiveness of *in situ* predator exposure as a method of improving the anti-predator traits of wildlife prey species should be evaluated. First, we must know if animals have learned or the traits of populations have changed after being exposed to a predator. This can be evaluated by comparing the anti-predator behaviour, physical and physiological traits and survival of predator-exposed versus predator naïve individuals. Second, we must know if these experiences with predators actually increase reintroduction success outside of predator-free sanctuaries. Detailed tracking of control versus predator-experienced animals and their progeny will be essential to determine the ultimate success of this strategy.

With proper monitoring, *in situ* predator exposure can be ethically defensible. Ultimately, *in situ* predator exposure has the potential to revolutionize reintroduction biology and significantly change the way that future faunal reintroductions are designed and implemented. Indeed, if the

goal is to restore ecosystems that contain both predators and their prey, such techniques may be essential.

Acknowledgements

Funding was provided by the Australian Research Council LP130100173. Frank Courchamp and the Blumstein lab commented on an earlier draft. Rebecca West provided valuable insight and contribution.

Literature Cited

- Allen, L., R. Engeman, and H. Krupa. 1996. Evaluation of three relative abundance indices for assessing dingo populations. *Wildlife Research* 23:197-205.
- Anson, J. R. and C. R. Dickman. 2013. Behavioral responses of native prey to disparate predators: naiveté and predator recognition. *Oecologia* 171:367-377.
- Armstrong, D. P., R. S. Davidson, W. J. Dimond, J. K. Perrott, I. Castro, J. G. Ewen, R. Griffiths, and J. Taylor. 2002. Population dynamics of reintroduced forest birds on New Zealand islands. *Journal of Biogeography* 29:609-621.
- Ashley, M. V., M. F. Willson, O. R. Pergams, D. J. O'Dowd, S. M. Gende, and J. S. Brown. 2003. Evolutionarily enlightened management. *Biological Conservation* 111:115-123.
- Banks, P. B. and C. R. Dickman. 2007. Alien predation and the effects of multiple levels of prey naiveté. *Trends in Ecology & Evolution* 22:229-230.
- Beck, B., L. Rapaport, M. S. Price, and A. C. Wilson. 1994. Reintroduction of captive-born animals. 265-286Book. *Reintroduction of captive-born animals*. Springer.

- Bekoff, M. 2002. The importance of ethics in conservation biology: Let's be ethicists not ostriches. *Endangered Species Update* 19:23-26.
- Berger, J., J. E. Swenson, and I.-L. Persson. 2001. Recolonizing carnivores and naive prey: conservation lessons from Pleistocene extinctions. *Science* 291:1036-1039.
- Biggins, D. E., A. Vargas, J. L. Godbey, and S. H. Anderson. 1999. Influence of prerelease experience on reintroduced black-footed ferrets (*Mustela nigripes*). *Biological Conservation* 89:121-129.
- Blumstein, D. T. 2006. The multipredator hypothesis and the evolutionary persistence of antipredator behavior. *Ethology* 112:209-217.
- Blumstein, D. T., J. C. Daniel, and B. P. Springett. 2004. A test of the multi-predator hypothesis: rapid loss of antipredator behavior after 130 years of isolation. *Ethology* 110:919-934.
- Blumstein, D. T., M. Mari, J. C. Daniel, J. G. Ardron, A. S. Griffin, and C. S. Evans. 2002. Olfactory predator recognition: wallabies may have to learn to be wary. *Animal Conservation* 5:87-93.
- Brown, C. and K. N. Laland. 2003. Social learning in fishes: a review. *Fish and Fisheries* 4:280-288.
- Burrows, N., D. Algar, A. Robinson, J. Sinagra, B. Ward, and G. Liddelow. 2003. Controlling introduced predators in the Gibson Desert of Western Australia. *Journal of Arid Environments* 55:691-713.
- Byers, J. A. 1997. American pronghorn: social adaptations and the ghosts of predators past. University of Chicago Press.

- Carroll, S. P., P. S. Jørgensen, M. T. Kinnison, C. T. Bergstrom, R. F. Denison, P. Gluckman, T. B. Smith, S. Y. Strauss, and B. E. Tabashnik. 2014. Applying evolutionary biology to address global challenges. *Science* 346:1245993.
- Carthey, A. J. and P. B. Banks. 2014. Naïveté in novel ecological interactions: lessons from theory and experimental evidence. *Biological Reviews* 84:932-949.
- Christensen, P. and N. Burrows. 1994. Project desert dreaming: experimental reintroduction of mammals to the Gibson Desert, Western Australia. In: *Reintroduction biology of Australian and New Zealand fauna*. (M. Serena ed.), pp. 199-207. Chipping Norton, NSW: Surrey Beatty & Sons.
- Clayton, J. A., C. R. Pavey, K. Vernes, and M. Tighe. 2014. Review and analysis of Australian macropod translocations 1969–2006. *Mammal Review* 44:109-123.
- Cooper, W. E., S. M. Diogo, and D. T. Blumstein. 2015. Fear, spontaneity, and artifact in economic escape theory: a review and prospectus. *Advances in the Study of Behavior* 47:147-179.
- Côté, I.M. and W.J. Sutherland. 1997. The effectiveness of removing predators to protect bird populations. *Conservation Biology* 11: 395–405.
- Côté, S. D., T. P. Rooney, J.-P. Tremblay, C. Dussault, and D. M. Waller. 2004. Ecological impacts of deer overabundance. *Annual Review of Ecology, Evolution, and Systematics* 35:113-147.
- Cox, J. G. and S. L. Lima. 2006. Naiveté and an aquatic–terrestrial dichotomy in the effects of introduced predators. *Trends in Ecology & Evolution* 21:674-680.

- Crisp, H. and K. Moseby. 2010. One-way gates: Initial trial of a potential tool for preventing overpopulation within fenced reserves. *Ecological Management & Restoration* 11:139-141.
- Crooks, K. R. and M. E. Soulé. 1999. Mesopredator release and avifaunal extinctions in a fragmented system. *Nature* 400:563-566.
- Darimont, C. T., S. M. Carlson, M. T. Kinnison, P. C. Paquet, T. E. Reimchen, and C. C. Wilmers. 2009. Human predators outpace other agents of trait change in the wild. *Proceedings of the National Academy of Sciences USA* 106:952-954.
- Ferrari, M. C., A. Gonzalo, F. Messier, and D. P. Chivers. 2007. Generalization of learned predator recognition: an experimental test and framework for future studies. *Proceedings of the Royal Society B: Biological Sciences* 274:1853-1859.
- Fischer, J., Lindenmayer, D. B. 2000. An assessment of the published results of animal relocations. *Biological Conservation* 96:1–11.
- Gaudioso, V. R., Sánchez-García, C., Pérez, J. A., Rodríguez, P. L., Armenteros, J. A., and M. E. Alonso, 2011. Does early antipredator training increase the suitability of captive red-legged partridges (*Alectoris rufa*) for releasing? *Poultry Science* 90:1900-1908.
- Gérard, A., H. Jourdan, C. Cugnière, A. Millon, and E. Vidal. 2014. Is naïveté forever? Alien predator and aggressor recognition by two endemic island reptiles. *Naturwissenschaften* 101:921-927.
- Gibson, D., G. Lundie-Jenkins, D. Langford, J. Cole, D. Clarke, and K. Johnson. 1994. Predation by feral cats, *Felis catus*, on the rufous hare-wallaby, *Lagorchestes hirsutus*, in the Tanami Desert. *Australian Mammalogy* 17:103-107.

- Griffin, A. 2004. Social learning about predators: a review and prospectus. *Animal Learning & Behavior* 32:131-140.
- Griffin, A. S., D. T. Blumstein, and C. S. Evans. 2000. Training captive-bred or translocated animals to avoid predators. *Conservation Biology* 14:1317-1326.
- Griffin, A. S. and C. S. Evans. 2003. The role of differential reinforcement in predator avoidance learning. *Behavioural Processes* 61:87-94.
- Hayward, M. W., L'Hotellier, F., O'Connor, T., Ward-Fear, G., Cathcart, J., Cathcart, T., Stephens, J., Stephens, J. Herman, K. and S. Legge. 2012. Reintroduction of bridled nailtail wallabies beyond fences at Scotia Sanctuary–Phase 1. *Proceedings of the Linnean Society of New South Wales* 134:A27-A37.
- Hayward, M. W. and G. I. Kerley. 2009. Fencing for conservation: restriction of evolutionary potential or a riposte to threatening processes? *Biological Conservation* 142:1-13.
- Hayward, M. W., G. I. Kerley, J. Adendorff, L. C. Moolman, J. O'brien, A. Sholto-Douglas, C. Bissett, P. Bean, A. Fogarty, and D. Howarth. 2007. The reintroduction of large carnivores to the Eastern Cape, South Africa: an assessment. *Oryx* 41:205-214.
- Hendry, A. P., T. J. Farrugia, and M. T. Kinnison. 2008. Human influences on rates of phenotypic change in wild animal populations. *Molecular Ecology* 17:20-29.
- Holden, C. and G. Mutze. 2003. Impact of rabbit haemorrhagic disease on introduced predators in the Flinders Ranges, South Australia. *Wildlife Research* 29:615-626.
- Hölzer, C., H. Bergmann, and I. McLean. 1995. Training captive-raised, naive birds to recognise their predator. *Research and Captive Propagation* (U. Ganslosser, JK Hodges & W. Kaumanns, eds). Filander Verlag, Fürth:198-206.
- Hunter, D.O., T. Britz, M. Jones and M. Letnic. 2015. Reintroduction of Tasmanian devils to

mainland Australia can restore top-down control in ecosystems where dingoes have been extirpated. *Biological Conservation* 191:428–435.

Innes, J., D. Kelly, J. M. Overton, and C. Gillies. 2010. Feathers to fur. *New Zealand Journal of Ecology* 34:86-114.

Islam, M., K. Ismail, and A. Boug. 2010. Catastrophic die-off of globally threatened Arabian oryx and sand gazelle in the fenced protected area of the arid central Saudi Arabia. *Journal of Threatened Taxa* 2:677-684.

Johnson, C. 2006. *Australia's Mammal Extinctions: A 50000 Year History*. Cambridge University Press.

King, C. M. 1984. *Immigrant Killers: Introduced Predators and the Conservation of Birds in New Zealand*. Oxford University Press Auckland.

Kohn, M. H., H.-J. Pelz, and R. K. Wayne. 2000. Natural selection mapping of the warfarin-resistance gene. *Proceedings of the National Academy of Sciences USA* 97:7911-7915.

Lahti, D. C., N. A. Johnson, B. C. Ajie, S. P. Otto, A. P. Hendry, D. T. Blumstein, R. G. Coss, K. Donohue, and S. A. Foster. 2009. Relaxed selection in the wild. *Trends in Ecology & Evolution* 24:487-496.

Letnic, M., E. G. Ritchie, and C. R. Dickman. 2012. Top predators as biodiversity regulators: the dingo *Canis lupus dingo* as a case study. *Biological Reviews* 87:390-413.

Lewis, J. C., K. L. Sallee, and R. T. Golightly Jr. 1999. Introduction and range expansion of nonnative red foxes (*Vulpes vulpes*) in California. *American Midland Naturalist* 142:372-381.

MacDonald, D. W. and L. A. Harrington. 2003. The American mink: the triumph and tragedy of adaptation out of context. *New Zealand Journal of Zoology* 30:421-441.

- Maloney, R. F. and I. G. McLean. 1995. Historical and experimental learned predator recognition in free-living New-Zealand robins. *Animal Behaviour* 50:1193-1201.
- Marlow, N.J., N.D. Thomas, A.A.E. Williams, B. Macmahon, J. Lawson, Y. Hitchen, J. Angus, and O. Berry. 2015. Cats (*Felis catus*) are more abundant and are the dominant predator of woylies (*Bettongia penicillata*) after sustained fox (*Vulpes vulpes*) control. *Australian Journal of Zoology* 16:18-27.
- McLean, I. G., C. Hölzer, and B. J. Studholme. 1999. Teaching predator-recognition to a naive bird: implications for management. *Biological Conservation* 87:123-130.
- McLean, I. G., G. Lundie-Jenkins, and P. J. Jarman. 1996. Teaching an endangered mammal to recognise predators. *Biological Conservation* 75:51-62.
- McLean, I. G., N. T. Schmitt, P. J. Jarman, C. Duncan, and C. Wynne. 2000. Learning for life: training marsupials to recognise introduced predators. *Behaviour* 137:1361-1376.
- Miller, B., D. Biggins, C. Wemmer, R. Powell, L. Calvo, L. Hanebury, and T. Wharton. 1990. Development of survival skills in captive-raised Siberian polecats (*Mustela eversmanni*) II: Predator avoidance. *Journal of Ethology* 8:95-104.
- Moseby, K. E., D.E.Peacock, and J.L. Read. 2015. Catastrophic cat predation: A call for predator profiling in wildlife protection programs. *Biological Conservation* 191:331-340.
- Moseby, K. E., A. Cameron, and H. A. Crisp. 2012. Can predator avoidance training improve reintroduction outcomes for the greater bilby in arid Australia? *Animal Behaviour* 83:1011-1021.
- Moseby, K. E. and B. M. Hill. 2011. The use of poison baits to control feral cats and red foxes in arid South Australia I. Aerial baiting trials. *Wildlife Research* 38:338-349.

- Moseby, K. E., J. L. Read, D. C. Paton, P. Copley, B. M. Hill, and H. A. Crisp. 2011. Predation determines the outcome of 10 reintroduction attempts in arid South Australia. *Biological Conservation* 144:2863-2872.
- Nogales, M., A. Martín, B. R. Tershy, C. Donlan, D. Veitch, N. Puerta, B. Wood, and J. Alonso. 2004. A review of feral cat eradication on islands. *Conservation Biology* 18:310-319.
- Phillips, B. L. and R. Shine. 2004. Adapting to an invasive species: toxic cane toads induce morphological change in Australian snakes. *Proceedings of the National Academy of Sciences USA* 101:17150-17155.
- Read, J. and Z. Bowen. 2001. Population dynamics, diet and aspects of the biology of feral cats and foxes in arid South Australia. *Wildlife Research* 28:195-203.
- Roy, S. S., C. G. Jones, and S. Harris. 2002. An ecological basis for control of the mongoose *Herpestes javanicus* in Mauritius: is eradication possible? In: *An ecological basis for control of the mongoose Herpestes javanicus in Mauritius: is eradication possible?* (C. R. Veitch and M. N. Clout eds), pp: 266-273. IUCN, Gland, Switzerland.
- Ruscoe, W. A., D. S. Ramsey, R. P. Pech, P. J. Sweetapple, I. Yockney, M. C. Barron, M. Perry, G. Nugent, R. Carran, and R. Warne. 2011. Unexpected consequences of control: competitive vs. predator release in a four - species assemblage of invasive mammals. *Ecology Letters* 14:1035-1042.
- Saunders, G. R., M. N. Gentle, and C. R. Dickman. 2010. The impacts and management of foxes *Vulpes vulpes* in Australia. *Mammal Review* 40:181-211.
- Savidge, J. A. 1987. Extinction of an island forest avifauna by an introduced snake. *Ecology* 68:660-668.

- Schakner, Z. and B. D.T. 2016. Learning and conservation behavior: an introduction and overview. In: Conservation Behavior: Applying behavioral ecology to wildlife conservation and management (O. Berger-Tal and D. Saltz, eds.), Cambridge University Press, Cambridge, XXX-YYY.
- Scofield, R. P., R. Cullen, and M. Wang. 2011. Are predator-proof fences the answer to New Zealand's terrestrial faunal biodiversity crisis? *New Zealand Journal of Ecology* 35:312-317.
- Sherley E.H, I.A.N. Stringer and G.R. Parish. 2010. *Summary of Native Bat, Reptile, Amphibian and Terrestrial Invertebrate Translocations in New Zealand*. Wellington: Department of Conservation
- Short, J. 2009. *The Characteristics and Success of Vertebrate Translocations within Australia*. Department of Agriculture, Fisheries and Forestry, Canberra, Australia.
- Short, J., J. Kinnear, and A. Robley. 2002. Surplus killing by introduced predators in Australia—evidence for ineffective anti-predator adaptations in native prey species? *Biological Conservation* 103:283-301.
- Short, J. and B. Turner. 2000. Reintroduction of the burrowing bettong *Bettongia lesueuri* (Marsupialia: Potoroidae) to mainland Australia. *Biological Conservation* 96:185-196.
- Sih, A., D. I. Bolnick, B. Luttbeg, J. L. Orrock, S. D. Peacor, L. M. Pintor, E. Preisser, J. S. Rehage, and J. R. Vonesh. 2010. Predator-prey naïveté, antipredator behavior, and the ecology of predator invasions. *Oikos* 119:610-621.
- Sinclair, A. R. E., R. P. Pech, C. R. Dickman, D. Hik, P. Mahon, and A. E. Newsome. 1998. Predicting effects of predation on conservation of endangered prey. *Conservation Biology* 12:564-575.

- Slotow, R., M. E. Garaï, B. Reilly, B. Page, and R. D. Carr. 2005. Population dynamics of elephants re-introduced to small fenced reserves in South Africa. *South African Journal of Wildlife Research* 35:23-32.
- Snyder, N. F., S. R. Derrickson, S. R. Beissinger, J. W. Wiley, T. B. Smith, W. D. Toone, and B. Miller. 1996. Limitations of captive breeding in endangered species recovery. *Conservation Biology* 10:338-348.
- Thompson, J. N. 1998. Rapid evolution as an ecological process. *Trends in Ecology & Evolution* 13:329-332.
- Towns, D. R., C. H. Daugherty, and A. Cree. 2001. Raising the prospects for a forgotten fauna: a review of 10 years of conservation effort for New Zealand reptiles. *Biological Conservation* 99:3-16.
- Towns, D. R. and S. M. Ferreira. 2001. Conservation of New Zealand lizards (Lacertilia: Scincidae) by translocation of small populations. *Biological Conservation* 98:211-222.
- van Heezik, Y., P. J. Seddon, and R. F. Maloney. 1999. Helping reintroduced houbara bustards avoid predation: effective anti-predator training and the predictive value of pre-release behaviour. *Animal Conservation* 2:155-163.
- Warburton, B. and K. Drew. 1994. Extent and nature of cyanide-shyness in some populations of Australian brushtail possums in New Zealand. *Wildlife Research* 21:599-605.
- Wayne, A. F., M. A. Maxwell, C. G. Ward, C. V. Vellios, I. Wilson, J. C. Wayne, and M. R. Williams. 2013. Sudden and rapid decline of the abundant marsupial *Bettongia penicillata* in Australia. *Oryx*:1-11.
- White Jr, T. H., J. A. Collazo, and F. J. Vilella. 2005. Survival of captive-reared Puerto Rican parrots released in the Caribbean National Forest. *The Condor* 107:424-432.

Williams, S. E. and E. A. Hoffman. 2009. Minimizing genetic adaptation in captive breeding programs: a review. *Biological Conservation* 142:2388-2400.

Wiseman, R., B. R. Page, and T. G. O'Connor. 2004. Woody vegetation change in response to browsing in Ithala Game Reserve, South Africa. *South African Journal of Wildlife Research* 34:25-37.

Wroe, S. and N. Milne. 2007. Convergence and remarkably consistent constraint in the evolution of carnivore skull shape. *Evolution* 61:1251-1260.