

# Are giant clams (*Tridacna maxima*) distractible? A multi-modal study

Ryan Doyle<sup>\*</sup>, Jonathan Kim<sup>\*</sup>, Angelika Pe<sup>\*</sup> and Daniel T. Blumstein

Department of Ecology and Evolutionary Biology, University of California, Los Angeles, Los Angeles, CA, United States of America

<sup>\*</sup>These authors contributed equally to this work.

## ABSTRACT

To properly assess risk, an animal must focus its attention on relevant external stimuli; however, attention can be reallocated when distracting stimuli are present. This reallocation of attention may interfere with an individual's ability to effectively assess risk and may impede its response. Multiple stimuli presented together can have additive effects as distractors, and these include stimuli in different modalities. Although changes in noise and water flow are detectable by some bivalves, this has not been studied in the context of risk assessment or distraction. We experimentally exposed giant clams (*Tridacna maxima*) to changes in water particle movement through underwater sound (motorboat noise) and increased water flow to determine whether these stimuli, individually or together, modified risk assessment or caused distraction. We found that clams responded to sound, flow, and their combination by increasing frequency of mantle retractions (a potential anti-predator response) when exposed to a stimulus. Sound alone did not change risk assessment in either the latency to close or to reemerge following closure. However, when exposed to both stimuli simultaneously, clams increased their latency to close. We suggest that clams perceive sound and flow in an additive way, and are thus distracted. Interestingly, and uniquely, clams discriminate these multimodal stimuli through a single sensory modality. For sessile clams, anthropogenic noise is detectable, yet unavoidable, suggesting that they be especially vulnerable to marine noise pollution.

Submitted 15 May 2020

Accepted 6 September 2020

Published 5 October 2020

Corresponding author

Daniel T. Blumstein,  
marmots@ucla.edu

Academic editor

James Reimer

Additional Information and  
Declarations can be found on  
page 11

DOI 10.7717/peerj.10050

© Copyright  
2020 Doyle et al.

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

**Subjects** Animal Behavior, Marine Biology, Zoology, Environmental Contamination and Remediation

**Keywords** Multimodal risk assessment, Giant clam, Multisensory risk assessment, Anthropogenic noise

## INTRODUCTION

Most animals experience some form of predation risk in their lifetimes (*Lima & Dill, 1990*). Individuals must be able to properly assess risk of predation, weighing the costs and benefits of either remaining or fleeing (*Caro, 2005*). Flight or other forms of protection are often associated with large energetic costs; individuals must expend energy to flee and lose out on potential energy that could have been obtained through feeding (*Ydenberg & Dill, 1986; Caro, 2005*).

Attention is the process of filtering out irrelevant stimuli from the environment in order for the individual to focus on biologically relevant stimuli (*Dukas, 2004*). Attention to tasks and external stimuli is limited, divisible and is sensitive to different modalities (*Dukas,*

2004; Chan et al., 2010; Blumstein, 2013). External stimuli can interfere with risk assessment by both masking biologically important signals (Zhou, Radford & Magrath, 2019) and distracting individuals, thus preventing them from focusing on avoiding predation (Rabin, Coss & Owings, 2006; Karp & Root, 2009; Chan et al., 2010). Distraction can also impair performance on other relevant tasks (Maes & Groot, 2003), potentially reducing feeding and reproductive opportunities. External acoustic and visual stimuli are commonly studied as distractions that impede anti-predatory responses in vertebrates (Grueninger & Pribram, 1969; Dukas, 2004; Wale, Simpson & Radford, 2013). However, there is less literature on distraction in invertebrates. A growing literature on giant clams' (*Tridacna maxima*) risk assessment shows that they respond to both visual and tactile threats (Stasek, 1965; Wilkens, 1986; Land, 2002; Soo & Todd, 2014; Johnson et al., 2016; Dehaudt et al., 2019).

However, bivalves assess risk in other modalities as well. They alter their anti-predator behaviors based on different chemical and auditory stimuli (Neo & Todd, 2011; Castorani & Hovel, 2015; Peng et al., 2016). They possess labial palps on their mantle, gills and siphon that can detect and respond to local water particle accelerations (Peng et al., 2016; Nedelec et al., 2016). Their palps also allow them to sense pressure changes from underwater sound waves and bivalves have been documented to significantly react to changes in underwater acoustic stimuli (e.g., Mosher, 1972; Peng et al., 2016; Shi et al., 2019; Charifi et al., 2018; Roberts et al., 2015).

Anthropogenic noise is one such external stimulus that could distract bivalves. Marine anthropogenic noise has been steadily rising over recent years due to shipping, motorboat activity, and pile-driving (Hildebrand, 2009). There remains an overall lack of study regarding the detrimental effects of noise with respect to invertebrates (Morley, Jones & Radford, 2014) and bivalves especially (Peng et al., 2016; Shi et al., 2019; Charifi et al., 2018; Roberts et al., 2015; Solan et al., 2016). Bivalves exposed to anthropogenic noise increase bioaccumulation of toxic metals (Peng et al., 2016), alter their digging, sediment uptake and valve closure patterns (Shi et al., 2019; Solan et al., 2016; Roberts et al., 2015) and experience reduced growth (Charifi et al., 2018). However, these studies did not look specifically at how noise potentially distracts bivalves from performing biologically important tasks. This research is critical because while mobile animals can respond to noise pollution through movement and migration away from areas of high intensity (Hirst & Rodhouse, 2000; Slabbekoorn et al., 2010), sessile marine invertebrates lack escape as an option. Since these species may have important roles in their community (e.g., Jordan & Valiela, 1982; Carballo & Naranjo, 2002; Neo et al., 2015; Solan et al., 2016), we need to know more about how they may be negatively impacted by anthropogenic stimuli. The little research that has been done calls for *in situ* testing and further understanding of the distracting effects of anthropogenic noise (Aguilar de Soto et al., 2013; Buscaino et al., 2016).

Acoustic stimuli propagate waves that oscillate water particles without creating a clear directional flow (Nedelec et al., 2016); however, these directional changes in underwater flow could also distract bivalves. The effect of changes in water flow has been investigated in bivalves in terms of how it influences their feeding (Levinton, 1991; Pilditch & Grant, 1999). However, we are aware of no studies on changes in flow as a potential bivalve distractor. Changes in water flow could indicate an increased potential for feeding; any increase in

underwater velocity could bring more edible plankton in the way of filter-feeding bivalves (Levinton, 1991) thereby altering bivalves' risk versus reward perception. Flow changes could also indicate changes in threat level because it is one of many factors naturally associated with the presence of a moving predator (McHenry et al., 2009).

Giant clams may commonly experience both stimuli in a multimodal way. Multimodal assessment may provide more information about threats and hence reduce uncertainty about risk leading to an increase or decrease in anti-predator responses (Munoz & Blumstein, 2012). Most multimodal research has explored the additive effect of multiple stimuli as a multisensory experience (Munoz & Blumstein, 2012; Munoz & Blumstein, 2019). Whereas a combination of sound and flow will indeed be considered multimodal, giant clams experience changes in water particle motion through a single sensory channel: their palps. If clams respond differently to multimodal stimulation compared to unimodal stimulation, then this will indicate that giant clams could modify their anti-predator behavior in a multimodal way and that this was achieved by a single sensory pathway binding multimodal stimuli. Specifically, the presence of additional stimuli in the form of sound or flow could potentially distract giant clams from the presence of a predator, causing a delay in how fast they retract their mantle.

We asked whether changes in water particle motion due to unimodal sound, unimodal water flow, or a multimodal combination of the two would distract giant clams or modify their risk assessment. When perturbed, giant clams will partially or fully retract their mantle as a potential anti-predatory response (Johnson et al., 2016; Dehault et al., 2019; Land, 2002; Wilkens, 1986). Thus, we used the latency to fully retract their mantle as an indicator of distraction levels and the latency for the mantle to fully re-emerge from its shell as a measure of perceived risk.

## MATERIALS & METHODS

### Study subjects

We marked giant clams ( $n = 32$ ) for subsequent study within Gump Reef in Cook's Bay, Mo'orea, French Polynesia ( $-17.482215, -149.827079$ ), a marine protected area that has been the site of previous studies of clam anti-predator behavior (Johnson et al., 2016; Dehault et al., 2019). Each clam was  $>5\text{m}$  apart from each other. We measured each clam's depth and the length of the each clam's shell. Clams received four total treatments over four separate days (with a planned day off between subsequent treatments) in a Latin square design: control, flow only, sound only, and flow and sound together. We initially planned to conduct all experiments over 8 days, but a large storm generated considerable runoff which filled the bay with sediment and created poor visibility making it impossible to study the clams. Thus, we waited until the water cleared and thus conducted two treatments 3 days later than planned. We refer to the first two treatments as pre-plume and the second two treatments as post-plume. Data were collected in the mornings, except for one day in which data had to be collected in the afternoon due to water clarity issues from storm water runoff. Clams were studied under permits issued by the Government of French Polynesia (permit approved on 21 November 2019).

### Flow calibration

We used an underwater aquarium pump (1/55 HP, ECO-FLO, Ashland OH) to manipulate water velocity over the clam. Before experimentation, we calibrated flow by video recording (Crosstour CT7000 underwater video camera, Shenzhen, China) the time it took red dye to travel 10 cm through a clear, plastic tube. We measured the velocity to be 0.43 ( $\pm 0.06$  SD) m/s ( $N = 10$  measurements) where the jet exiting the nozzle of the pump was 15 cm away from the opening of the tube.

### Sound calibration

We used an Oceanears DRS-8 MOD 2 underwater speaker (North Canton, OH) to broadcast recordings of outboard motorboats collected underwater. Five recordings of motorboat sounds of similar amplitude were obtained from a previous study ([Simpson et al., 2016](#)). The five motorboat recordings consist of a research vessel (5-m-long aluminum hull boat with 30 hp Suzuki outboard motors) motoring 10–200 m away from the hydrophone at various speeds, mimicking boat operations that are common in coral reef environments ([Simpson et al., 2016](#)).

To determine how the broadcast sounds compared to that of the reef environment, we broadcast and re-recorded our exemplars underwater using a Wildlife Acoustics Song Meter SM2+ (Concord, MA) with a Wildlife Acoustics hydrophone. We first calibrated the hydrophone by playing 45 s of white noise of a known amplitude (90 dB re 20  $\mu$ Pa measured at 1 m) through an Oceanears DRS-8 MOD 2 underwater speaker (North Canton, OH) above water. The white noise was generated in Audacity (version 2.3.3, 2019) and was measured with a Radioshack (33–2055) Digital Sound Level Meter set to weighting A (Radioshack, Fort Worth, Texas) placed 1 m away from the speaker. The white noise was then broadcast underwater 1 m from the hydrophone at 0.5 m depth. We also similarly broadcasted the five motorboat exemplars (89–91 dB re 20  $\mu$ Pa measured at 1 m) against the calibrated hydrophone. In Praat (version 6.1.109, 2020), we calculated the average intensity (minimum pitch = 100 Hz) of the white noise and then adjusted the scale to reflect the intensity measured above water to which we added 61.5 dB to roughly estimate intensity re 1  $\mu$ Pa underwater ([National Research Council, 1994](#); [Gausland, 2000](#)). This created a scaling factor that we used to adjust the amplitude of each motorboat exemplar. Using Praat, we calculated the average intensity of our stimuli across exemplars, which was 150.6 dB ( $\pm 1.03$  SD) re 1  $\mu$ Pa.

### Stimulus presentation

Clams received four different treatments over four days: control (pump and speaker off), flow only (pump on, speaker off), sound only (pump off, speaker on) and flow and sound (pump and speaker on) in a Latin Square design. For all four treatments, the clam was subjected to the presence of both the pump and speaker, whether on or off, for 60 s before it was perturbed. The pump and speaker were attached to a 1.5 m PVC pipe. The hose from the pump was positioned 15 cm away from the clam while the speaker was 1 m away pointed at the center of the clam's mantle. This ensured that the clam was subjected to 0.4 ( $\pm 0.07$  SD) m/s of flow and 149.13 dB ( $\pm 1.03$  SD) re 1  $\mu$ Pa. Clams receiving sound or flow and sound treatment were randomly exposed to one of five motor boat noises.

During this 60 s period before perturbation, we counted the number of mantle twitches (partial retractions) as an indicator of perceived threat. The entire length of the clam's mantle was then rubbed back and forth twice using a pencil's eraser attached to a different 2 m long PVC pipe that permitted the person to be  $\geq 1$  m away. The latency to close was recorded using either Crosstour CT7000 or a GoPro Hero 4 camera. The latency to re-open was measured both in the field using a digital stopwatch, and using the camera. We defined latency to close as the first sign of retraction of the mantle to the point where the clam is fully or almost closed. We defined latency to reemerge as the time from the first contact of the pencil to the clam returning to its previous undisturbed state that the observer has determined prior to the clam being rubbed. If the clam did not fully reemerge within 3 min, we determined reemergence as the point where its mantle was most fully extended. The camera recorded the entire process from the 60 s before perturbation until the observer was confident the clam had fully reemerged. In 16 cases we were unable to confidently measure the latency to re-open in the field. Due to the extremely high correlation between field and camera playback latency times, we estimated these recorded times from field observations using a prediction equation based on the correlation between measurements (estimated time recorded from video =  $1.0263 * \text{stopwatch time} + 1.9489$ ;  $R^2 = 0.98$ ;  $p < 0.001$ ,  $N = 103$  measurements to estimate latency to re-emerge).

To reduce experimental interference, we did not record the precise velocity of ambient water flow during the experiments. However, we noted flow as either negligible or noticeable by observing pieces of macroalgae moving underwater after conducting each experimental treatment. If pieces of macroalgae in the water column at the same depth of the clam moved in a single direction, flow was marked as noticeable. We estimated an upper limit range of ambient velocity during testing by recording the velocity at which a piece of macroalgae moved 10 cm through a clear, plastic tube with a GoPro nine separate times during a day with particularly high underwater flow. We estimated this velocity as  $0.09 (\pm 0.064 \text{ SD})$  m/s, which was substantially lower than our experimental velocity. Thus, we infer that our experimental velocity was substantially greater than background changes in flow.

### Statistical analysis

Prior to analysis, we plotted histograms of variables to check for normality and  $\log_{10}$  transformed retractions and latency to re-emerge.

To determine if stimuli were detected by giant clams, we fitted a linear mixed model with the number of retractions as the dependent variable using the lme4 version 1.1-21, (Bates et al., 2014) and lmerTest version 3.1-1 (Kuznetsova, Brockhoff & Christensen, 2017) packages in R version 3.6.2 (R Core Team, 2019). Treatment, trial number, clam size, and plume presence (defined as before or after the disruptive plume) were added as fixed effects. To account for repeated measures, subject ID was added as a random effect. We tested the assumptions of our mixed model by plotting a histogram of residuals (they were approximately normal), plotting a q-q plot (they were approximately linear), and plotting fitted versus predicted values (there was no obvious pattern). We also compared our model to a linear model without subject ID as a random effect to determine if there was a significant difference in individuals' number of retractions. We estimated marginal

means using the MuMIn version 1.43.15 (Barton, 2019) and emmeans version 1.4.4 (Lenth et al., 2020) packages in R and estimated the d-score as a measure of treatment effect size. We plotted estimated marginal means using plotly version 4.9.2 (Sievert et al., 2020).

To determine if multimodal stimuli distracted giant clams, we fitted a linear mixed model as above with the latency to close, and another model with the latency to re-emerge as dependent variables. Again, we added treatment, trial number, clam size, and plume presence as fixed effects and added subject ID as a random effect. Model fit was determined by examining the normality of residuals in a histogram and quantile plot. Again, both models were compared to a linear model without the random effect of subject ID. Estimated marginal means were plotted with standard errors.

We tested for exemplar effects by creating a dataset containing only those where motorboat noise was used and then fitting each of our models with sound exemplar as a fixed factor to determine if any variation in response was explained by the specific exemplar used.

All data and code are contained in Tables S1, S2. Throughout, we set our alpha to 0.05, and thus interpret  $p < 0.05$  as significant.

## RESULTS

After controlling for significant random effects explained by individual (Chi-square = 2.83, 1 df,  $p = 0.007$ ), and non-significant fixed effects of trial number, plume presence and clam size, there were more twitches recorded during a 60 s period when clams were exposed to sound, flow and the combination of the two stimuli compared to the no stimuli treatment (Table 1A, Linear mixed effects model,  $p < 0.05$  for all). There were large effect sizes between the control and each treatment (Fig. 1A, Table S3;  $d > 0.75$ ). Additionally, there were moderate effect sizes ( $0.75 > d > 0.4$ ) between unimodal sound and unimodal flow as well as between unimodal sound and multimodal flow and sound (Fig. 1A, Table S3).

After controlling for significant random effects explained by individual (Chi-square = 10.848, 1 df,  $p < 0.001$ ), significant fixed effects of clam size, and non-significant fixed effects of trial number and plume presence, we found that the multimodal treatment (combination of sound and flow) significantly increased clams' latency to close (Table 1B, Linear mixed effects model,  $p < 0.05$ ) while the other two unimodal treatments (flow only and sound only) had no effect on latency to close when compared to the control (Table 1B, Linear mixed effects model,  $p > 0.05$ ). There was a small to moderate effect size between multimodal flow and sound and the control; all other comparisons had small effect sizes (Fig. 1B, Table 1B;  $d = 0.35$  and  $d < 0.3$ , respectively).

After controlling for significant random effects explained by individual (Chi-square = 22.442, 1 df,  $p < 0.001$ ), significant fixed effects of plume presence and non-significant fixed effects of trial number and clam size, we found that clams did not significantly change their latency to reemerge after treatments when compared to the control (Table 1C, Linear mixed effects model,  $p > 0.05$  for all). We found moderate effect sizes ( $0.75 > d > 0.4$ ) between unimodal flow treatment compared to all other treatments as well as the control.


When we tested for exemplar effects, variation in sound exemplars did not explain any variation in clam response.

**Table 1** Results from linear mixed effects model on variation in clam response. This model explains variation in giant clam responses (a, number of retractions in 60 s, b, latency to close, c, latency to reemerge) following a multimodal stimulus experiment. The reference level was no stimulus.

Fixed effects	Estimate $\pm$ SE	P
(a) Number of retractions in 60 s		
Intercept	0.535 $\pm$ 0.144	<0.001
Trial Number	0.0198 $\pm$ 0.347	0.569
After Plume Presence	-0.111 $\pm$ 0.077	0.150
Size	-0.323 $\pm$ 1.18	0.787
Treatment		
Flow	0.280 $\pm$ 0.05	<0.001
Sound	0.169 $\pm$ 0.05	0.001
Flow + Sound	0.277 $\pm$ 0.05	<0.001
Subject ID	0.010	0.101
(b) Latency to close		
Intercept	2.490 $\pm$ 1.125	0.032
Trial Number	-0.081 $\pm$ 0.248	0.743
After Plume Presence	0.188 $\pm$ 0.550	0.733
Size	19.685 $\pm$ 9.437	0.046
Treatment		
Flow	0.177 $\pm$ 0.348	0.613
Sound	0.054 $\pm$ 0.361	0.882
Flow + Sound	0.749 $\pm$ 0.353	0.037
Subject ID	0.753	0.867
(c) Latency to reemerge		
Intercept	1.465 $\pm$ 0.271	<0.001
Trial Number	0.012 $\pm$ 0.051	0.816
After Plume Presence	0.368 $\pm$ 0.112	0.002
Size	0.331 $\pm$ 2.346	0.889
Treatment		
Flow	-0.114 $\pm$ 0.074	0.127
Sound	0.014 $\pm$ 0.079	0.859
Flow + Sound	0.024 $\pm$ 0.074	0.752
Subject ID	0.056	0.236

## DISCUSSION

Animals often use cues from a variety of modalities to reduce uncertainty about their environment (Dall & Johnstone, 2002), which potentially permits more precise assessments of risk. Here we used two stimuli, from different modalities, either alone or together, to study anti-predator responses in clams. We found that clams increased their frequency of partial retractions during the 60 s acclimation period after exposure to any treatment. Clam mantle retraction is presumably a discrete anti-predator response used for protection and to scare potential predators with the water expelled from its siphon during retraction (Wilkins, 1986; Land, 2002).



**Figure 1** A comparison of estimated marginal means of (A)  $\text{Log}_{10}$  (number of partial retractions) (B) latency to close (C)  $\text{Log}_{10}$  (Latency to reemerge) between all treatments. Bars indicate one standard error from the means. Letters above bars indicate treatments that are significantly different from each other.

Full-size [DOI: 10.7717/peerj.10050/fig-1](https://doi.org/10.7717/peerj.10050/fig-1)



While twitches were not complete valve closures, the twitches were still a physical response that is closely linked to the species' highest level antipredator response—full mantle closure. Increased anti-predator behavior indicates that motorboat noise may act as a giant clam stressor (*Wright et al., 2007*). Previous studies have shown the adverse effects of anthropogenic noise on bivalves' physiological processes such as metabolism and growth rate (*Roberts et al., 2015; Peng et al., 2016; Charifi et al., 2018; Shi et al., 2019*). Thus, our results add to previous studies in assessing whether the affected physiological processes in bivalves are the result of anthropogenic noise-based distraction.

Sound and flow did not significantly alter clams' latency to close when tested as separate unimodal treatments. However, when sound and flow were presented together as a multimodal stimulus, clams closed more slowly, indicating that the presentation of multiple stimuli can be distracting. Given that there was no significant effect of each stimulus when presented on its own and that giant clams perceive both stimuli through their palps, a unisensory receptor, we conclude that these stimuli have an additive effect on compromising clams' attention. These results are consistent with the distracted prey hypothesis in which peripheral stimuli may compromise an individual's attention from biologically necessary tasks (*Chan et al., 2010*). Less vigilant prey become more likely targets for predators (*Krause & Godin, 1996*), and a slower closing time in clams encountering a multimodal stimulus indicates that distracted clams may be more vulnerable to predatory damage.

Clams did not differ in their reemergence time between exposure to treatment and control. However, the flow as a unimodal treatment had a modest, but not significant, decrease in reemergence time from the control (*Table 1*) and had a moderate to large effect size when compared to unimodal sound and multimodal sound and flow. This modest decrease in reemergence time in response to increased flow around the clam may indicate an increased opportunity to filter feed because there are more sediments and nutrients moving through its environment (*Taghon, Nowell & Jumars, 1980; Trager, Hwang & Strickler, 1990; Skilleter & Peterson, 1994; Hardy & Hardy, 1969; Hawkins & Klumpp, 1995*). Thus, clams may reemerge faster to take advantage of increased nutrient availability. As a result, increased water flow cannot be considered a neutral distractor. It is rather unique in that it is a biologically significant stimulus, forcing clams to factor foraging opportunity into their decision-making process. Clams must simultaneously manage predation risk and obtain energy through feeding behavior: two potentially exclusive behaviors with their own set of benefits and consequences. If clams reemerge sooner, they could potentially make the mistake of exposing themselves to predators; however, if they stay closed for too long, they are missing out on feeding opportunities. Further studies are needed that explore reemergence time of clams in areas with naturally high levels of flow to determine if they behave differently from clams in areas, like our study site on Gump Reef, with relatively low flow.

These findings further illustrate the negative effects that anthropogenic noise can have on bivalves (*Peng et al., 2016; Shi et al., 2019; Charifi et al., 2018; Roberts et al., 2015*). These prior studies identified alterations in essential processes such as digging behavior, metabolism, and growth in bivalves (*Peng et al., 2016; Shi et al., 2019; Charifi et al., 2018*).

Of the prior studies that identified negative effects, none, until this one, specifically focused on anthropogenic noise as a distraction. A recent meta-analysis reported anthropogenic noise negatively affects families across the animal kingdom, potentially altering distribution, communication, foraging, and homeostasis in these families (Kunc & Schmidt, 2019). However, species' responses differ widely based on ecological context and physical constraints (Wright et al., 2007). While mobile species can avoid anthropogenic noise stressors (e.g., Francis, Ortega & Cruz, 2009; Francis et al., 2011; Goodwin & Shriver, 2011; Nowacek et al., 2007; Weilgart, 2007; Weilgart, 2018), giant clams are, for the most part, sessile and must endure any anthropogenic disturbance. Giant clams play important ecological roles on coral reefs, including water filtration and nutrient sequestration (Neo et al., 2015). Thus, anything that makes more clams more vulnerable to predators may contribute to the loss of these important services that giant clams provide.

Multimodal stimuli are traditionally defined as separate stimuli that are perceived through different sensory channels (Partan & Marler, 2005). Many animals are able to detect multiple stimuli across different modalities and create a unified percept by integrating them (Brown & Magnavacca, 2003; Partan et al., 2010). In contrast to all other studies we are aware of, giant clams perceive multimodal stimuli through a single sensory modality, their palps, but nevertheless integrated these stimuli. It is important to note that giant clams depend heavily on visual cues for risk assessment (Wilkins, 1986; Stasek, 1965). Our manipulations did not create any shading events or cause any visible turbulence, thus we infer that clams were not relying on visual cues to sense the particle motion caused by noise or flow. We therefore suggest that local changes in water flow and changes in underwater noise are indeed multimodal although perceived through a single sensory channel by clams. The clams' ability to react differently to either unimodal or multimodal treatments of sound and flow is a novel example of how animals can bind multimodal stimulus together through a single sensory pathway to modify their risk assessment.

Sound and flow changes sensed by clams could also be viewed as multi-attribute stimuli. Unlike multimodal stimuli, which are typically perceived through different sensory channels, multi-attribute stimuli are perceived through a single sensory channel. Multi-attribute stimuli are often studied in bioacoustics (e.g., Gerhardt, 1992; Castellano & Rosso, 2007; Naguib & Wiley, 2001; Gerhardt & Schul, 1999; Kershenbaum et al., 2016). In these cases, different components of an acoustic signal, including rate and frequency rises, can send multiple messages or further inform an individual. Previous studies on crayfish (*Orconectes virilis*) demonstrated that animals can use multiple separate chemical cues perceived through a single olfactory channel (Hazlett, 1999; Bouwma & Hazlett, 2001). Female Italian treefrogs (*Hyla intermedia*) weight separate components of male mating calls differently when selecting a mate (Castellano & Rosso, 2007). Similarly, great tits (*Parus major*) assess multiple acoustic attributes of song (Weary, 1990). Female bluehead wrasse (*Thalassoma bifasciatum*) use multiple visual cues, such as coloration and courtship display, to assess risk (Warner & Dill, 2000). In honeybees, different olfactory stimuli are processed through different clusters of glomeruli, enabling them to process different components of odor through different pathways (Abel, Rybak & Menzel, 2001; Kirschner et al., 2006). The majority of these previous studies have focused on multi-attribute signaling and not strictly

multi-attribute risk assessment (*Munoz & Blumstein, 2012; Munoz & Blumstein, 2019*). In addition, these previous studies focused mainly on terrestrial organisms; more studies are needed on multi-attribute stimulus assessment in marine organisms. We have shown that giant clams can bind together multiple stimuli into one sensory channel, an ability that we suspect is shared with other species.

## CONCLUSIONS

Giant clams increased their frequency of mantle retractions, a potential anti-predator behavior, in response to motorboat playback, increased flow and the combination of the two. Clams also slowed their closing times when exposed to a combination of sound and flow, indicating distraction through multimodal means. These findings add evidence to the deleterious nature of anthropogenic noise on sessile marine invertebrates and present a novel case of multimodal stimuli processed together through a single sensory channel. We recommend further study of unisensory systems and their relationships to traditionally multimodal stimuli.

## ACKNOWLEDGEMENTS

We thank the Richard Gump South Pacific Research Station for logistical support, Zachary Schakner for the playback equipment, Maud Ferrari for motorboat recordings, John Milligan and Vivian Kim for help with our equipment, and Dana Williams for logistical and statistical help as well as comments on a previous version of this article. The article was improved by addressing the comments of three astute reviewers.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This work was supported by the UCLA Department of Ecology and Evolutionary Biology. There was no additional external funding received for this study. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:  
UCLA Department of Ecology and Evolutionary Biology.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Ryan Doyle, Jonathan Kim and Angelika Pe conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Daniel T. Blumstein conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.

## Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

This research was conducted under a permit issued by the Government of French Polynesia (permit approved on 21 November 2019).

## Data Availability

The following information was supplied regarding data availability:

Data and code are available at Github.

The data sheets are available at:

[https://github.com/distractedclams/Clam-Distraktion/blob/master/Clam\\_Multimodal\\_Datasheet\\_FINAL.csv](https://github.com/distractedclams/Clam-Distraktion/blob/master/Clam_Multimodal_Datasheet_FINAL.csv).

The R Code is available at:

[https://github.com/distractedclams/Clam-Distraktion/blob/master/Distracted\\_Clams\\_FINAL.R](https://github.com/distractedclams/Clam-Distraktion/blob/master/Distracted_Clams_FINAL.R).

The Experimental Trial Video is available at: <https://www.youtube.com/watch?v=O8GkVb8yeMo>

The Experimental Design is available at: <https://github.com/distractedclams/Clam-Distraktion/blob/master/Presentation.pdf>.

## Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.10050#supplemental-information>.

## REFERENCES

- Abel R, Rybak J, Menzel R. 2001. Structure and response patterns of olfactory interneurons in the honeybee, *Apis mellifera*. *Journal of Comparative Neurology* **437**:363–383 DOI [10.1002/cne.1289](https://doi.org/10.1002/cne.1289).
- Aguilar de Soto NA, Delorme N, Atkins J, Howard S, Williams J, Johnson M. 2013. Anthropogenic noise causes body malformations and delays development in marine larvae. *Science Reports* **3**:2831 DOI [10.1038/srep02831](https://doi.org/10.1038/srep02831).
- Barton K. 2019. MuMIn: multi-model inference. R package version, 1.43.6. Available at <https://CRAN.R-project.org/package=MuMIn>.
- Bates D, Maechler M, Bolker B, Walker S. 2014. Linear mixed-effects models using ‘eigen’ and S4. *Journal of Statistical Software* **67**:1–48.
- Blumstein DT. 2013. Attention, habituation, and antipredator behaviour: implications for urban birds. In: Gil D, Brumm H, eds. *Avian urban ecology: behavioural and physiological adaptations*. Oxford: Oxford University Press, 41–53.
- Bouwma P, Hazlett BA. 2001. Integration of multiple predator cues by the crayfish *Orconectes propinquus*. *Animal Behaviour* **61**:771–776 DOI [10.1006/anbe.2000.1649](https://doi.org/10.1006/anbe.2000.1649).
- Brown GE, Magnavacca G. 2003. Predator inspection behaviour in a characin fish: an interaction between chemical and visual information? *Ethology* **109**:739–750 DOI [10.1046/j.1439-0310.2003.00919.x](https://doi.org/10.1046/j.1439-0310.2003.00919.x).

- Buscaino G, Ceraulo M, Pieretti N, Corrias V, Farina A, Filiciotto F, Maccarrone V, Grammatta R, Caruso F, Giuseppe A, Mazzola S. 2016.** Temporal patterns in the soundscape of the shallow waters of a Mediterranean marine protected area. *Science Report* **6**:34230 DOI [10.1038/srep34230](https://doi.org/10.1038/srep34230).
- Carballo JL, Naranjo S. 2002.** Environmental assessment of a large industrial marine complex based on a community of benthic filter-feeders. *Marine Pollution Bulletin* **44**:605–610 DOI [10.1016/S0025-326X\(01\)00295-8](https://doi.org/10.1016/S0025-326X(01)00295-8).
- Caro T. 2005.** *Antipredator defenses in birds and mammals*. Chicago: University of Chicago Press.
- Castellano S, Rosso A. 2007.** Female preferences for multiple attributes in the acoustic signals of the Italian treefrog, *Hyla intermedia*. *Behavioral Ecology and Sociobiology* **61**:1293–1302 DOI [10.1007/s00265-007-0360-z](https://doi.org/10.1007/s00265-007-0360-z).
- Castorani MCN, Hovel KA. 2015.** Native predator chemical cues induce anti-predation behaviors in an invasive marine bivalve. *Biological Invasions* **18**:169–181.
- Chan AAYH, Giraldo-Perez P, Smith S, Blumstein DT. 2010.** Anthropogenic noise affects risk assessment and attention: the distracted prey hypothesis. *Biology Letters* **6**:458–461 DOI [10.1098/rsbl.2009.1081](https://doi.org/10.1098/rsbl.2009.1081).
- Charifi M, Miserazzi A, Sow M, Perrigault M, Gonzalez P, Ciret P, Benomar S, Massabuau J. 2018.** Noise pollution limits metal bioaccumulation and growth rate in a filter feeder, the Pacific oyster *Magallana gigas*. *PLOS ONE* **13**:e0194174 DOI [10.1371/journal.pone.0194174](https://doi.org/10.1371/journal.pone.0194174).
- Dall SRX, Johnstone RA. 2002.** Managing uncertainty: information and insurance under the risk of starvation. *Philosophical Transactions of the Royal Society B Biological Sciences* **357**:1519–1526 DOI [10.1098/rstb.2002.1061](https://doi.org/10.1098/rstb.2002.1061).
- Dehaut B, Nguyen M, Vadlamudi A, Blumstein DT. 2019.** Giant clams discriminate threats along a risk gradient and display varying habituation rates to different stimuli. *Ethology* **125**:392–398 DOI [10.1111/eth.12863](https://doi.org/10.1111/eth.12863).
- Dukas R. 2004.** Causes and consequences of limited attention. *Brain, Behavior and Evolution* **63**:197–210 DOI [10.1159/000076781](https://doi.org/10.1159/000076781).
- Francis CD, Ortega CP, Cruz A. 2009.** Noise pollution changes avian communities and species interactions. *Current Biology* **19**:1415–1419 DOI [10.1016/j.cub.2009.06.052](https://doi.org/10.1016/j.cub.2009.06.052).
- Francis CD, Paritsis J, Ortega CP, Cruz A. 2011.** Landscape patterns of avian habitat use and nest success are affected by chronic gas well compressor noise. *Landscape Ecology* **26**:1269–1280 DOI [10.1007/s10980-011-9609-z](https://doi.org/10.1007/s10980-011-9609-z).
- Gausland I. 2000.** Impact of seismic surveys on marine life. *The Leading Edge* **19**:903–905 DOI [10.1190/1.1438746](https://doi.org/10.1190/1.1438746).
- Gerhardt HC. 1992.** Multiple messages in acoustic signals. *Seminars in Neuroscience* **4**:391–400 DOI [10.1016/1044-5765\(92\)90047-6](https://doi.org/10.1016/1044-5765(92)90047-6).
- Gerhardt HC, Schul J. 1999.** A quantitative analysis of behavioral selectivity for pulse rise-time in the gray treefrog, *Hyla versicolor*. *Journal of Comparative Physiology A* **185**:33–40 DOI [10.1007/s003590050363](https://doi.org/10.1007/s003590050363).
- Goodwin SE, Shriver WG. 2011.** Effects of traffic noise on occupancy patterns of forest birds. *Conservation Biology* **25**:406–411.

- Grueninger WE, Pribram KH. 1969.** Effects of spatial and nonspatial distractors on performance latency of monkeys with frontal lesions. *Journal of Comparative and Physiological Psychology* **68**:203–209 DOI [10.1037/h0027498](https://doi.org/10.1037/h0027498).
- Hardy JT, Hardy SA. 1969.** Ecology of *Tridacna* in Palau. *Pacific Science* **23**:467–472.
- Hawkins AJS, Klumpp DW. 1995.** Nutrition of the giant clam *Tridacna gigas* (L.). II Relative contributions of filter-feeding and the ammonium-nitrogen acquired and recycled by symbiotic alga towards total nitrogen requirements for tissue growth and metabolism. *Journal of Experimental Marine Biology and Ecology* **190**:263–290 DOI [10.1016/0022-0981\(95\)00044-R](https://doi.org/10.1016/0022-0981(95)00044-R).
- Hazlett BA. 1999.** Responses to multiple chemical cues by the crayfish *Orconectes virilis*. *Behaviour* **136**:161–171 DOI [10.1163/156853999501261](https://doi.org/10.1163/156853999501261).
- Hildebrand JA. 2009.** Anthropogenic and natural sources of ambient noise in the ocean. *Marine Ecology Progress Series* **395**:5–20 DOI [10.3354/meps08353](https://doi.org/10.3354/meps08353).
- Hirst AG, Rodhouse PG. 2000.** Impacts of geophysical seismic surveying on fishing success. *Reviews in Fish Biology and Fisheries* **10**:113–118 DOI [10.1023/A:1008987014736](https://doi.org/10.1023/A:1008987014736).
- Johnson GC, Karajah MT, Mayo K, Armenta TC, Blumstein DT. 2016.** The bigger they are the better they taste: size predicts predation risk and anti-predator behavior in giant clams. *Journal of Zoology* **301**:102–107.
- Jordan TE, Valiela I. 1982.** A nitrogen budget of the ribbed mussel, *Geukensia demissa*, and its significance in nitrogen flow in a New England salt marsh. *Limnology and Oceanography* **27**:75–90 DOI [10.4319/lo.1982.27.1.0075](https://doi.org/10.4319/lo.1982.27.1.0075).
- Karp DS, Root TL. 2009.** Sound the stressor: How hoatzins (*Opisthocomus hoazin*) react to ecotourist conversation. *Biodiversity and Conservation* **18**:3733–3742 DOI [10.1007/s10531-009-9675-6](https://doi.org/10.1007/s10531-009-9675-6).
- Kershenbaum A, Blumstein DT, Roch MA, Acay C, Backus G, Bee MA, Bohn K, Cao Y, Carter C, Casar C, Coen M, DeRuiter SL, Doyle L, Edelman S, Ferrer-i Cancho R, Freeberg TM, Garland EC, Gustison M, Harley HE, Huetz C, Hughes M, Bruno JH, Ilany A, Jin DZ, Johnson M, Ju C, Karnowski J, Lohr B, Manser MB, McCowan B, Mercado III E, Narins PM, Piel A, Rice M, Salmi R, Sasahara K, Sayigh L, Shiu Y, Taylor C, Vallejo EE, Waller S, Zamora-Guiterrez V. 2016.** Acoustic sequences in non-human animals: a tutorial review and prospectus. *Biological Reviews* **91**:13–52 DOI [10.1111/brv.12160](https://doi.org/10.1111/brv.12160).
- Kirschner S, Kleineidam CJ, Zube C, Rybak J, Grunewald B, Rossler W. 2006.** Dual olfactory pathway in the honeybee, *Apis mellifera*. *Journal of Comparative Neurology* **499**:933–952 DOI [10.1002/cne.21158](https://doi.org/10.1002/cne.21158).
- Krause J, Godin JGJ. 1996.** Influence of prey foraging posture on flight behavior and predation risk: predators take advantage of unwary prey. *Behavioural Ecology* **7**:264–271 DOI [10.1093/beheco/7.3.264](https://doi.org/10.1093/beheco/7.3.264).
- Kunc HP, Schmidt R. 2019.** The effects of anthropogenic noise on animals: a meta-analysis. *Biology Letters* **15**:Article 20190649 DOI [10.1098/rsbl.2019.0649](https://doi.org/10.1098/rsbl.2019.0649).
- Kuznetsova A, Brockhoff PB, Christensen RHB. 2017.** Tests in linear mixed effects models. *Journal of Statistical Software* **82**:1–26.

- Land MF. 2002.** The spatial resolution of the pinhole eyes of giant clams (*Tridacna maxima*). *Proceedings of the Royal Society B: Biological Sciences* **270**:185–188.
- Lenth R, Singmann H, Love J, Buerkner P, Herve M. 2020.** Estimated marginal means, aka least-squares means. Available at <https://CRAN.Rproject.org/package=emmeans>.
- Levinton JS. 1991.** Variable feeding behavior in three species of *Macoma* (Bivalvia: Tellinacea) as a response to water flow and sediment transport. *Marine Biology* **110**:375–383 DOI [10.1007/BF01344356](https://doi.org/10.1007/BF01344356).
- Lima SL, Dill LM. 1990.** Behavioral decisions made under the risk of predation: a review and prospectus. *Canadian Journal of Zoology* **68**:619–640 DOI [10.1139/z90-092](https://doi.org/10.1139/z90-092).
- Maes JHR, De Groot G. 2003.** Effects of noise on the performance of rats in an operant discrimination task. *Behavioural Processes* **61**:57–68 DOI [10.1016/S0376-6357\(02\)00163-8](https://doi.org/10.1016/S0376-6357(02)00163-8).
- McHenry MJ, Feitl KE, Strother JA, Van Trump WJ. 2009.** Larval zebrafish rapidly sense the water flow of a predator's strike. *Biology Letters* **5**:477–479 DOI [10.1098/rsbl.2009.0048](https://doi.org/10.1098/rsbl.2009.0048).
- Morley EL, Jones G, Radford AN. 2014.** The importance of invertebrates when considering the impacts of anthropogenic noise. *Proceedings of the Royal Society B: Biological Sciences* **281**:20132683 DOI [10.1098/rspb.2013.2683](https://doi.org/10.1098/rspb.2013.2683).
- Mosher JJ. 1972.** The responses of *Macoma balthica* (Bivalvia) to vibrations. *Journal of Molluscan Studies* **40**:125–131.
- Munoz NE, Blumstein DT. 2012.** Multisensory perception in uncertain environments. *Behavioural Ecology* **23**:457–462 DOI [10.1093/beheco/arr220](https://doi.org/10.1093/beheco/arr220).
- Munoz NE, Blumstein DT. 2019.** Optimal multisensory integration. *Behavioural Ecology* **23**:457–462.
- Naguib M, Wiley RH. 2001.** Review: estimating the distance to a source of sound: mechanisms and adaptations for long-range communication. *Animal Behavior* **62**:825–837 DOI [10.1006/anbe.2001.1860](https://doi.org/10.1006/anbe.2001.1860).
- National Research Council. 1994.** *Low-frequency sound and marine mammals: current knowledge and research needs*. Washington, D.C.: The National Academies Press.
- Nedelec SL, Campbell J, Radford AN, Simpson SD, Merchant ND. 2016.** Particle motion: the missing link in underwater acoustic ecology. *Methods in Ecology and Evolution* **7**:836–842 DOI [10.1111/2041-210X.12544](https://doi.org/10.1111/2041-210X.12544).
- Neo ML, Eckman W, Vicentuan K, Teo SLM, Todd PA. 2015.** The ecological significance of giant clams in coral reef ecosystems. *Biological Conservation* **181**:111–123 DOI [10.1016/j.biocon.2014.11.004](https://doi.org/10.1016/j.biocon.2014.11.004).
- Neo ML, Todd PA. 2011.** Predator-induced changes in fluted giant clam (*Tridacna squamosa*) shell morphology. *Journal of Experimental Marine Biology and Ecology* **397**:21–26 DOI [10.1016/j.jembe.2010.11.008](https://doi.org/10.1016/j.jembe.2010.11.008).
- Nowacek DP, Thorne LH, Johnston DW, Tyack PL. 2007.** Responses of cetaceans to anthropogenic noise. *Mammal Review* **37**:81–115 DOI [10.1111/j.1365-2907.2007.00104.x](https://doi.org/10.1111/j.1365-2907.2007.00104.x).
- Partan SR, Fulmer AG, Gounard MAM, Redmond JE. 2010.** Multimodal alarm behavior in urban and rural gray squirrels studied by means of observation and a mechanical robot. *Current Zoology* **56**:313–326 DOI [10.1093/czoolo/56.3.313](https://doi.org/10.1093/czoolo/56.3.313).

- Partan SR, Marler P. 2005.** Issues in the classification of multimodal communication signals. *The American Naturalist* **166**:231–245 DOI [10.1086/431246](https://doi.org/10.1086/431246).
- Peng C, Zhao X, Liu S, Shi W, Han Y, Guo C, Jiang J, Wan H, Shen T, Guangxu L. 2016.** Effects of anthropogenic sound on digging behavior, metabolism, Ca<sup>2+</sup>/Mg<sup>2+</sup> ATPase activity, and metabolism-related gene expression of the bivalve *Sinonovacula constricta*. *Science Reports* **6**:24266 DOI [10.1038/srep24266](https://doi.org/10.1038/srep24266).
- Pilditch CA, Grant J. 1999.** Effect of variations in flow velocity and phytoplankton concentration on sea scallop (*Placopecten magellanicus*) grazing rates. *Journal of Experimental Marine Biology and Ecology* **240**:111–136 DOI [10.1016/S0022-0981\(99\)00052-0](https://doi.org/10.1016/S0022-0981(99)00052-0).
- R Core Team. 2019.** R: a language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. Available at <https://www.R-project.org/>.
- Rabin LA, Coss RG, Owings DH. 2006.** The effects of wind turbines on antipredator behavior in California ground squirrels (*Spermophilus beecheyi*). *Biological Conservation* **131**:410–420 DOI [10.1016/j.biocon.2006.02.016](https://doi.org/10.1016/j.biocon.2006.02.016).
- Roberts L, Cheesman S, Breithaupt T, Elliott M. 2015.** Sensitivity of the mussel *Mytilus edulis* to substrate-borne vibration in relation to anthropogenically generated noise. *Marine Ecology Progress Series* **538**:185–195 DOI [10.3354/meps11468](https://doi.org/10.3354/meps11468).
- Shi W, Han Y, Guan X, Rong J, Du X, Zha S, Tang Y, Liu G. 2019.** Anthropogenic noise aggravates the toxicity of cadmium on some physiological characteristics of the blood clam *Tegillarca granosa*. *Frontiers in Physiology* **10**:Article 3389.
- Sievert C, Parmer C, Hocking T, Chamberlain S, Ram K, Corvellec M, Despouy P. 2020.** Create Interactive WebGraphics via ‘plotly.js’. Available at <https://CRAN.R-project.org/package=plotly>.
- Simpson SD, Radford AN, Nedelec SL, Ferrari MCO, Chivers DP, McCormick MI, Meekan MG. 2016.** Anthropogenic noise increases fish mortality by predation. *Nature Communications* **7**:Article 10544 DOI [10.1038/ncomms10544](https://doi.org/10.1038/ncomms10544).
- Skilleter GA, Peterson CH. 1994.** Control of foraging behavior of individuals within an ecosystem context: the clam *Macoma balthica* and interactions between competition and siphon cropping. *Oecologia* **100**:268–278 DOI [10.1007/BF00316954](https://doi.org/10.1007/BF00316954).
- Slabbekoorn H, Bouton N, Van Opzeeland I, Coers A, Ten Cate C, Popper AN. 2010.** A noisy spring: the impact of globally rising underwater sound levels on fish. *Trends in Ecology and Evolution* **25**:419–427 DOI [10.1016/j.tree.2010.04.005](https://doi.org/10.1016/j.tree.2010.04.005).
- Solan M, Hauton C, Godbold JA, Wood CL, Leighton TG, White P. 2016.** Anthropogenic sources of underwater sound can modify how sediment-dwelling invertebrates mediate ecosystem properties. *Scientific Reports* **6**:20540 DOI [10.1038/srep20540](https://doi.org/10.1038/srep20540).
- Soo P, Todd PA. 2014.** The behaviour of giant clams (Bivalvia: Cardiidae: Tridacninae). *Marine Biology* **161**:2699–2717 DOI [10.1007/s00227-014-2545-0](https://doi.org/10.1007/s00227-014-2545-0).
- Stasek CR. 1965.** Behavioural adaptation of the giant clam *Tridacna maxima* to the presence of grazing fishes. *Veilger* **8**:29–35.
- Taghon GL, Nowell ARM, Jumars PA. 1980.** Induction of suspension feeding in spionid polychaetes by high particulate fluxes. *Science* **210**:562–564 DOI [10.1126/science.210.4469.562](https://doi.org/10.1126/science.210.4469.562).



- Trager GC, Hwang JS, Strickler JR. 1990.** Barnacle suspension-feeding in variable flow. *Marine Biology* **105**:117–127 DOI [10.1007/BF01344277](https://doi.org/10.1007/BF01344277).
- Wale MA, Simpson SD, Radford AN. 2013.** Noise negatively affects foraging and antipredator behaviour in shore crabs. *Animal Behavior* **86**:111–118 DOI [10.1016/j.anbehav.2013.05.001](https://doi.org/10.1016/j.anbehav.2013.05.001).
- Warner RR, Dill LM. 2000.** Courtship displays and coloration as indicators of safety rather than of male quality: the safety assurance hypothesis. *Behavioural Ecology* **11**:444–451 DOI [10.1093/beheco/11.4.444](https://doi.org/10.1093/beheco/11.4.444).
- Weary DM. 1990.** Categorization of song notes in great tits: which acoustic features are used and why? *Animal Behavior* **39**:450–457 DOI [10.1016/S0003-3472\(05\)80408-7](https://doi.org/10.1016/S0003-3472(05)80408-7).
- Weilgart LS. 2007.** A brief review of known effects of noise on marine mammals. *International Journal of Comparative Psychology* **20**:159–168.
- Weilgart LS. 2018.** The impact of ocean noise pollution on fish and invertebrates. Report for OceanCare, Switzerland.
- Wilkins LA. 1986.** The visual system of the giant clam *Tridacna*: behavioral adaptations. *Biological Bulletin* **170**:393–408 DOI [10.2307/1541850](https://doi.org/10.2307/1541850).
- Wright AJ, Hatch LT, Aguilar de Soto N, Baldwin AL, Bateson M, Beale CM, Clark C, Deak T, Edwards EF, Fernandez A, Godinho A, Hatch LT, Kakuschke A, Lusseau D, Martineau D, Romero ML, Weilgart LS, Wintle BA, Notarbartolo-di Sciara G, Cortes Martin V. 2007.** Anthropogenic noise as a stressor in animals: a multidisciplinary perspective. *Journal of Comparative Psychology* **20**:250–273.
- Ydenberg RC, Dill LM. 1986.** The economics of fleeing from predators. *Advances in the Study of Behavior* **16**:229–249 DOI [10.1016/S0065-3454\(08\)60192-8](https://doi.org/10.1016/S0065-3454(08)60192-8).
- Zhou Y, Radford AN, Magrath RD. 2019.** Why does noise reduce response to alarm calls? Experimental assessment of masking, distraction and greater vigilance in wild birds. *Functional Ecology* **33**:1280–1289 DOI [10.1111/1365-2435.13333](https://doi.org/10.1111/1365-2435.13333).